

# Endoscopy Auckland

Please ensure you have returned your Patient Particulars form.

## PREPARATION FOR COLONOSCOPY AND GASTROSCOPY

### YOUR APPOINTMENT IS:

Date: \_\_\_\_\_ Day: \_\_\_\_\_ Arrival time: \_\_\_\_\_

***Please adhere to instructions given by Endoscopy Auckland instead of instructions given with MOVIPREP.***

A colonoscopy is a visual examination of the bowel using a narrow flexible telescope. To obtain good views of the bowel lining, the bowel must be completely empty. This is achieved with the help of a preparation called **MOVIPREP - you will need to contact your chemist at least 4 days prior to your appointment to ensure they have the MOVIPREP available for you.** (You will need a box of MOVIPREP).

The gastroscopy is also a visual examination using a narrow flexible telescope to view the stomach and its lining. If you suffer from any medical condition that may require specialised information or treatment during preparation/procedure, please contact Endoscopy Auckland for advice.

### MEDICATIONS:

If you are taking Iron tablets please discontinue them one (1) week prior to the test. If you are taking Aspirin, Cartia, Warfarin or any other anti-coagulant tablets regularly, please advise Endoscopy Auckland beforehand. If you are a diabetic it is important to notify Endoscopy Auckland, as there are special instructions for preparation. You can take all other usual medications 1 hour prior to beginning or 1 hour after completing MOVIPREP to allow for absorption.

### THINGS TO DO PRIOR TO YOUR APPOINTMENT

You are advised to contact your medical insurance company to obtain a prior approval number or letter. This makes payment easier.

Organise a family member or friend (or a taxi) to drop you at Endoscopy Auckland and collect you approximately 1½ hours after your appointment time.

**Because a sedative is used it is very important you DO NOT drive or operate machinery or dangerous appliances, on the day of the procedure.**

### THINGS TO BRING WITH YOU TO YOUR APPOINTMENT

Any x-rays and/or barium studies relevant to your examination (if available).  
Your medications or a list of them.  
Your medical insurance prior approval number or letter (if relevant).

### ACCOUNTS ARE PAYABLE AT THE COMPLETION OF THE PROCEDURE (if no prior approval has been obtained)

If for some reason you also require antibiotics (you will be told at the time) there is an extra cost of between \$5.00 - \$55.00.

**THREE DAYS BEFORE EXAMINATION until after the examination, avoid foods containing seeds, nuts, pips, wholegrains.**

### DAY BEFORE EXAMINATION

- **Have a light breakfast** and a **light lunch** (see over for light diet suggestions).
- Do not be tempted to eat a heavy, solid meal.
- **After lunch STOP ALL FOOD until after your procedure** – drink only *clear fluids*.

### At least 1 hour after lunch

- Mix together the contents of 1 sachet A and 1 sachet B in a jug, then add in a litre of water. Stir the solution until the powder has dissolved and the MOVIPREP solution is clear or slightly hazy. Drink the entire litre over an hour. Then over the next ½ hour, drink 2 x 250ml glassfuls of *clear fluid*. Allow another TWO hours for the MOVIPREP to work. You will experience increasingly watery stools, this is normal, so ensure you can get to a toilet easily.

### 3 hours after the first dose

- Mix together the contents of the remaining sachet A and sachet B in a jug, then add in a litre of water. Stir the solution until the powder has dissolved and the MOVIPREP solution is clear or slightly hazy. Drink the entire litre over an hour. Follow this with 2 x 250ml glassfuls of *clear fluid* over the next ½ hour. Allow another TWO hours for MOVIPREP to work. You will experience increasingly watery stools, this is normal, so ensure you can get to a toilet easily. When you are no longer experiencing watery stools go to bed as usual.

***Clear fluids*** - water, tea/coffee, clear soup, soft drink, cordial, fruit juice without pulp. **NO BLACKCURRANT OR ANY RED OR PURPLE DRINKS.**

### DAY OF EXAMINATION – STOP ALL FLUIDS FROM MIDNIGHT

- **FOR A MORNING APPOINTMENT:**  
**STOP ALL FLUIDS FROM MIDNIGHT**
- **FOR AN AFTERNOON APPOINTMENT:**  
**STOP ALL FLUIDS SIX HOURS BEFORE APPOINTMENT**

**Note:** Because this preparation will cause multiple watery bowel motions it is advisable that you remain within easy reach of toilet facilities, therefore you may be more comfortable in your home environment. Individual responses to laxatives do vary - the preparation often works within 30 minutes but may take up to three hours.

**ONLY BY CARRYING OUT THE PREPARATION AS ABOVE WILL THE BOWEL BE CLEAN ENOUGH FOR THE PROCEDURE TO BE CARRIED OUT.**

Do not hesitate to telephone **Endoscopy Auckland** if you have any queries or doubts about this appointment or about the preparation.

## SUGGESTED DIET BEFORE COLONOSCOPY

### LIGHT DIET

Rice bubbles, or cornflakes (use reduced fat milk only)  
Eggs (not fried), omelettes, plain tinned spaghetti  
White bread or toast (not high fibre)  
Plain crackers (not wholemeal or multigrain), plain scones (no sultanas, dates), plain biscuits (Arrowroot or Wine)  
Butter, margarine, cheese  
Honey, marmalade  
Marmite, Vegemite  
Fish, skinless chicken (not fried or highly seasoned)  
White rice, plain pasta, skinless potatoes or kumara

**NO fruit, vegetables, pips, seeds, red meat, brown bread, muesli.**

### CLEAR FLUIDS

*You are encouraged to drink as much clear fluid as possible to help clear your bowel and prevent dehydration. For comfort or to give variation of taste, try alternating clear fluids during your bowel preparation.*

Water  
Lemon flavoured energy drinks, colourless cordials and carbonated drinks  
Strained fruit juice, some soups (e.g. strained chicken noodle)  
Lemon ice blocks, light coloured jellies, barley sugars, boiled sweets  
Tea and coffee – you may use a little reduced fat milk if you wish but do not consume milk based products like latte, cappuccino or yoghurt

**NO highly coloured (i.e. red, purple, green) drinks.**

### PLEASE NOTE:

If you are feeling nauseated, gingerale may help due to its natural anti-nausea properties.

It is not unusual to feel cold or bloated when drinking the bowel preparation. Wear warm comfortable clothing and try short walks to help alleviate bloating. Vaseline can be wiped around the anus to reduce discomfort during bowel cleansing. It is important to drink **the full amount** of bowel preparation to achieve adequate cleansing of the bowel.

*Please contact Endoscopy Auckland if you require further assistance.*




**ENDOSCOPY**

**AUCKLAND**

## COLONOSCOPY AND GASTROSCOPY

### Instructions using MOVIPREP



148 GILLIES AVENUE EPSOM AUCKLAND 1023 NEW ZEALAND  
PH: 09 623 2020 0800 ENDOSCOPY 0800 363 672 FAX: 09 623 1010